The Bryologist 92(4), 1989, pp. 558–570
Copyright © 1989 by the American Bryological and Lichenological Society, Inc.

Index to Volume 92 of THE BRYOLOGIST[1,2]

Abietinella, 162
Acarospora cervina var. glaucocarpa, 108; fuscata, 108, 275; immersa, 275; sinopica, 407+
Acaulon, 411; muticum var. rufescens, new to Alabama, 485
Achrophyllum, 262+
Acrocladium, 159
Acrocordia gemmata, 275; megalospora, 275
Acrophyllum dentatum, 266; quadrifarium, 266
Actinodontium, 268; standleyi, 266
Actinothuidium hookeri, 121+
Adelanthus brevicaulis, *346*; decipiens, 330+
Adelotheciaceae, 262+
Adelothecium, systematic position of, 261+; bogotense, 261+, 264*, 463
Africa, Lejeunea capensis, disjunct with South America, 305
AHMADJIAN, V. [Review of:] D. H. Dalby, D. L. Hawksworth & S. L. Jury (eds.), Horizons in Lichenology, 150
AKIYAMA, H. A reevaluation of Hypnodendron opacum (Hypnodendraceae, Musci), 198
Alabama, Acaulon muticum var. rufescens new to, 485; Aphanorrhegma serratum new to, 485; Astomum muhlenbergianum new to, 486; Bruchia brevifolia new to, 130; Campylostelium saxicola new to, 482; Didymodon tophaceus new to, 486; Ephemerum spinulosum new to, 486; Eucladium verticillatum new to, 487; Fontinalis sullivantii new to, 483; Fontinalis welchiana new to, 483; Helodium paludosum new to, 487; Herpetineuron toccoae new to, 483; Isopterygium distichaceum new to, 483; Luisierella barbula new to, 487; Micromitrium austinii new to, 487; M. megalosporum new to, 487; Platydictya confervoides new to, 487; Pleuridium ravenelii new to, 487; Polytrichum pallidisetum new to, 483; Schizomitrium pallidum new to, 487; Splachnobryum obtusum new to, 488; Tortula rhizophylla new to, 483; Trichostomum crispulum new to, 483; new and rare mosses from, 482+, 485
Alaska, microhabitat preferences of Dactylina in, 467+; nitrogen fixation of lichens in, 202
Alberta, saxicolous lichen flora in, 105
Alectoria nigricans, 501; ochroleuca, 501
AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN. Arachidonic and eicosapentaenoic acids in lipids of Bryum bicolor Dicks. Effects of controlled temperature and illumination, 178
ALI, A. M. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 178
ALLEN, B. H. [Review of:] J. E. Beever, A Key to the Genera of New Zealand Mosses, 450
ALLEN, B. H. [Review of:] G. R. Brassard (ed.), Bryophyta Exsiccata Terrae-Novae et Labradoricae, 152
ALLEN, B. H. [Review of:] S. P. Churchill & P. M. Eckel (eds.), Musci Planitiebus Incolae Exsiccati, Mosses of the Great Plains of North America, 152
ALLEN, B. H. [Review of:] H. Inoue (ed.), Bryophyta Selecta Exsiccata, 448
ALLEN, B. H. [Review of:] C. M. Matteri (ed.), Musci Fuegiani Exsiccati, Serie I. Nos. 76–100, 555
ALLEN, B. H. *See* MAGILL, R. E. & B. H. ALLEN, 434
ALLEN, B. *See* WHITTEMORE, A. & B. ALLEN, 261
ALLEN, B. H. & R. E. MAGILL. Recent literature on mosses—142, 543
Alnus, 202, 203; incana, 206
Alobiella husnotii, 330+
Alobiellopsis dominicensis, 330+
Aloina aloides, 309; bifrons, 359; brevirostris, 359; rigida, 359
Amblyodon dealbatus, 369+*
Amblyotropis, 268+; setosa, 266
Amblystegium 159; humile, *533*; leitense, 129; riparium, 196; serpens, 485; tenax, 196+, 498; tenax var. spinifolium, 485
Amphidium mougeotii, 482
Amygdalaria, 25+, 57; elegantior, 56+; panaeola, 57
Anacamptodon splachnoides, 482
Anacolia, 373+; abyssinica, *372*; campylopus, *372*; intertexta, 372; laevisphaera, 369+*; setifolia, *372*; sinensis, *372*
Anaptychia palmulata, 275
Anastrophyllum hellerianum, 350+; minutum, 500
ANDERSON, L. E. *See* CRUM, H. & L. E. ANDERSON, 533
ANDERSON, L. E. *See* SHAW, J., B. D. MISHLER & L. E. ANDERSON, 314
Andes, epiphyte in montane forests of, 216
ANDO, H., T. SEKI & W. B. SCHOFIELD. Generic distinctness of Brotherella from Pylaisiadelpha (Musci), 209
Andoa, 159
Andreaea megistospora subsp. epapillosa, *533*; m. var. **epapillosa** stat. nov., 533; rupestris var. papillosa, 473+
Aneura pinguis, 331+, 500
Anisomeridium biforme, 275
Anomobryum filiforme, 113+, 463; steerei, 418
Anomoclada mucosa, *344*; **portoricensis** comb. nov., 344+
Anomodon, 122, 381+; abbreviatus, 381; acutifolius, *385*; attenuatus, 381+, 463; dentatus, 381+*; fragillimus, *383+*; longifolius, 383; minor, 121; perarmatus, *383+*; rostratus, 383, 463; tasmanicus, *383+*; thraustus, 381; viticulosus, 381+; wrightii, 383
Anoplolejeunea conferta, 330+
Anthoceros, 219; carolinianus, *346*; communis, 344; kuntzeanus, 344; punctatus, 331+
Anthocerotae, of Puerto Rico and the Virgin Islands, 329+
Anthoxanthum odoratum, 73
antiherbivore function, lichen secondary compounds in, 326+
antimicrobial function, lichen secondary compounds in, 326+
Aongstroemia jamaicensis, 462; longipes, 473+; orientalis, 463
Aphanolejeunea crenata, 330+; diaphana, 346; exigua, 330+; sicaefolia, 330+; sintenisii, 346
Aphanorrhegma serratum, new to Alabama, 485
Apotomanthus succulentus, *346*
Aptychella proligera, 462

[1] Compiled by Robert E. Magill, Department of Bryology, Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299.

[2] New scientific names and combinations are in boldface; page numbers of names in synonomy are in italic; taxa mentioned several times in a paper are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

0007-2745/89/558–570$1.45/0

The Bryologist 92(4), 1989, pp. 558–570
Copyright © 1989 by the American Bryological and Lichenological Society, Inc.

Index to Volume 92 of THE BRYOLOGIST[1,2]

Abietinella, 162
Acarospora cervina var. glaucocarpa, 108; fuscata, 108, 275; immersa, 275; sinopica, 407+
Acaulon, 411; muticum var. rufescens, new to Alabama, 485
Achrophyllum, 262+
Acrocladium, 159
Acrocordia gemmata, 275; megalospora, 275
Acrophyllum dentatum, 266; quadrifarium, 266
Actinodontium, 268; standleyi, 266
Actinothuidium hookeri, 121+
Adelanthus brevicaulis, *346*; decipiens, 330+
Adelotheciaceae, 262+
Adelothecium, systematic position of, 261+; bogotense, 261+, 264*, 463
Africa, Lejeunea capensis, disjunct with South America, 305
AHMADJIAN, V. [Review of:] D. H. Dalby, D. L. Hawksworth & S. L. Jury (eds.), Horizons in Lichenology, 150
AKIYAMA, H. A reevaluation of Hypnodendron opacum (Hypnodendraceae, Musci), 198
Alabama, Acaulon muticum var. rufescens new to, 485; Aphanorrhegma serratum new to, 485; Astomum muhlenbergianum new to, 486; Bruchia brevifolia new to, 130; Campylostelium saxicola new to, 482; Didymodon tophaceus new to, 486; Ephemerum spinulosum new to, 486; Eucladium verticillatum new to, 487; Fontinalis sullivantii new to, 483; Fontinalis welchiana new to, 483; Helodium paludosum new to, 487; Herpetineuron toccoae new to, 483; Isopterygium distichaceum new to, 483; Luisierella barbula new to, 487; Micromitrium austinii new to, 487; M. megalosporum new to, 487; Platydictya confervoides new to, 487; Pleuridium ravenelii new to, 487; Polytrichum pallidisetum new to, 483; Schizomitrium pallidum new to, 487; Splachnobryum obtusum new to, 488; Tortula rhizophylla new to, 483; Trichostomum crispulum new to, 483; new and rare mosses from, 482+, 485
Alaska, microhabitat preferences of Dactylina in, 467+; nitrogen fixation of lichens in, 202
Alberta, saxicolous lichen flora in, 105
Alectoria nigricans, 501; ochroleuca, 501
AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN. Arachidonic and eicosapentaenoic acids in lipids of Bryum bicolor Dicks. Effects of controlled temperature and illumination, 178
ALI, A. M. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 178
ALLEN, B. H. [Review of:] J. E. Beever, A Key to the Genera of New Zealand Mosses, 450
ALLEN, B. H. [Review of:] G. R. Brassard (ed.), Bryophyta Exsiccata Terrae-Novae et Labradoricae, 152
ALLEN, B. H. [Review of:] S. P. Churchill & P. M. Eckel (eds.), Musci Planitiebus Incolae Exsiccati, Mosses of the Great Plains of North America, 152
ALLEN, B. H. [Review of:] H. Inoue (ed.), Bryophyta Selecta Exsiccata, 448
ALLEN, B. H. [Review of:] C. M. Matteri (ed.), Musci Fuegiani Exsiccati, Serie I. Nos. 76–100, 555
ALLEN, B. H. *See* MAGILL, R. E. & B. H. ALLEN, 434
ALLEN, B. *See* WHITTEMORE, A. & B. ALLEN, 261
ALLEN, B. H. & R. E. MAGILL. Recent literature on mosses—142, 543
Alnus, 202, 203; incana, 206
Alobiella husnotii, 330+
Alobiellopsis dominicensis, 330+
Aloina aloides, 309; bifrons, 359; brevirostris, 359; rigida, 359
Amblyodon dealbatus, 369+*
Amblyotropis, 268+; setosa, 266
Amblystegium 159; humile, *533*; leitense, 129; riparium, 196; serpens, 485; tenax, 196+, 498; tenax var. spinifolium, 485
Amphidium mougeotii, 482
Amygdalaria, 25+, 57; elegantior, 56+; panaeola, 57
Anacamptodon splachnoides, 482
Anacolia, 373+; abyssinica, *372*; campylopus, *372*; intertexta, 372; laevisphaera, 369+*; setifolia, *372*; sinensis, *372*
Anaptychia palmulata, 275
Anastrophyllum hellerianum, 350+; minutum, 500
ANDERSON, L. E. *See* CRUM, H. & L. E. ANDERSON, 533
ANDERSON, L. E. *See* SHAW, J., B. D. MISHLER & L. E. ANDERSON, 314
Andes, epiphyte in montane forests of, 216
ANDO, H., T. SEKI & W. B. SCHOFIELD. Generic distinctness of Brotherella from Pylaisiadelpha (Musci), 209
Andoa, 159
Andreaea megistospora subsp. epapillosa, *533*; m. var. **epapillosa** stat. nov., 533; rupestris var. papillosa, 473+
Aneura pinguis, 331+, 500
Anisomeridium biforme, 275
Anomobryum filiforme, 113+, 463; steerei, 418
Anomoclada mucosa, *344*; **portoricensis** comb. nov., 344+
Anomodon, 122, 381+; abbreviatus, 381; acutifolius, *385*; attenuatus, 381+, 463; dentatus, 381+*; fragillimus, *383+*; longifolius, 383; minor, 121; perarmatus, *383+*; rostratus, 383, 463; tasmanicus, *383+*; thraustus, 381; viticulosus, 381+; wrightii, 383
Anoplolejeunea conferta, 330+
Anthoceros, 219; carolinianus, *346*; communis, 344; kuntzeanus, 344; punctatus, 331+
Anthocerotae, of Puerto Rico and the Virgin Islands, 329+
Anthoxanthum odoratum, 73
antiherbivore function, lichen secondary compounds in, 326+
antimicrobial function, lichen secondary compounds in, 326+
Aongstroemia jamaicensis, 462; longipes, 473+; orientalis, 463
Aphanolejeunea crenata, 330+; diaphana, 346; exigua, 330+; sicaefolia, 330+; sintenisii, 346
Aphanorrhegma serratum, new to Alabama, 485
Apotomanthus succulentus, *346*
Aptychella proligera, 462

[1] Compiled by Robert E. Magill, Department of Bryology, Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299.

[2] New scientific names and combinations are in boldface; page numbers of names in synonomy are in italic; taxa mentioned several times in a paper are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

0007-2745/89/558–570$1.45/0

arachidonic acid, in lipids of Bryum bicolor, 178+
Arachniopsis coactilis, *346*; diacantha, 330+
Araucaria araucana, 217
Archidium alternifolium, 485; tenerrimum, 485
Archilejeunea parviflora, 330+; viridissima, *346*
Arctoparmelia centrifuga, 107+; subcentrifuga, 107+
Arctostaphylos nummularia, 126
Argentina, additions to the moss flora of Tierra del Fuego, 174
Arnellia fennica, 500
Arthonia caesia, 275+; punctiformis, 275; radiata, 275, 455
Arthopyrenia punctiformis, 275
Arthothelium spectabile, 275
Arthrocormus, 493
Arthrorhaphis alpina, 108
Arundinaria falcata, 120
Asahinea chrysantha, 501
Aschisma kansanum, 411
Asia, two new species of Syrrhopodon from southeast, 186
Aspicilia, 50; caesiocinerea, 108; cinerea, 107+; gibbosa, 326+; subsorediza, *68*+
Aspiromitus, 219
Asterella elegans, 331+
Asterothyrium rotuliforme, 68+
ASTHANA, A. K. *See* SRIVASTAVA, S. C. & A. K. ASTHANA, 219
Astomiopsis amblyocalyx, 225+*; exserta, 225+*; × **altivallis** sp. nov., 227+*
Astomum ludovicianum, 486; muhlenbergianum, new to Alabama, 486
Astrophyllum curvatulum, *533*
Athalamia hyalina, 360
Atrichum, 323; crispum, 474; oerstedianum, 462
Aulacomniaceae, 368+
Aulacomnium acuminatum, 498; androgynum, 369+*; heterostichum, 314+, 369+*; palustre, 95+, 369+*, 498; turgidum, 498
Australia, Pleurophascum occidentale new to, 299; the genus Schlotheimia in, 282+; two new taxa of Syrrhopodon from, 302
Austrocedrus chilensis, 217
axillary hairs, of Bartramiaceae, 369+

Bacidia apiahica, 68; atrogrisea, *68*+; chlorantha, 455; endoleuca, *68*+; inundata, 275; laurocerasi, 68+, 275; medialis, 68; naegelii, 275; polychroa, 275; rubella, 275; sabuletorum, 275; schweinitzii, 275; sphaeroides, *68*+, 275; trachona, 275
Baeomyces roseus, 275+; rufus, 278
Barbella cubensis, 463
Barbilophozia atlantica, 500; attenuata, 350+, 500; barbata, 500; binsteadii, 500; hatcheri, 500; kunzeana, 500; lycopodioides, 500; quadriloba, 500
Barbula indica, 463; obtusula, *533*; platyneura, 356+, 361; spiralis, 462; steerei, 418
Bartramia, 371+; sect. Acoleos, orth. var., 374+; sect. Acoleus, 373+; sect. Breutelia, 373+; sect. Lycopodiobryum, 374+; sect. Polyptychium, 373+; subsect. Anacoliopsis, 373+; abyssinica, *372*; affinis, 373; afroaffinis, nom. nud., 373; angustifolia, 369+*; breutelii, 373; campylopus, *372*; chrysura, 373+; comiramea, *374*; commutata, *373*; comosa, 374, *376*; crassa, 374; crassicaulis, 374; cygnea, *376*; erecta, *376*; fendleri, *376*; globosa, *376*; grandis, 374; intertexta, 372; lycopodium, nom. nud., 374; microstoma, 462; patula, *376*; polygastrica, 373; setifolia, *372*; tuerckheimii, *376*
Bartramiaceae, axillary hairs of, 369+; taxonomic and phylogenetic studies on, 368; subfam. Bartramioideae, 371; subfam. **Breutelioideae** subfam. nov., 373+; subfam. **Conostomoideae** subfam. nov., 372+
Bartramidula, 372+, 374+; bartramioides, 375+; cernua, 374, *376*; comosa, 375+, *376*; dispersa, *376*; erecta, 374+; globosa, 374+; le-testui, *376*; mexicana, *376*; nana, *376*; patula, 375; pusilla, *376*; roylei, 375; setifolia, *372*; treubii, 375+; tuerckheimii, 375+; wilsonii, 374
BASILE, D. V., B. K. KUSHNER & M. R. BASILE. A new method for separating and comparing arabinogalactan proteins for the chemosystematics of the Hepaticae, 164
BASILE, M. R. *See* BASILE, D. V., B. K. KUSHNER & M. R. BASILE, 164
Bazzania bidens, 330; breuteliana, 330+*; cubensis, 330+; cuneistipula, 330+; eggersiana, 344; falcata, 330+; gracilis, 330+; hookeria, 330+; longa, 330+; longistipula, 330+; pycnophylla, 330+; roraimensis, 344; schwaneckeana, 330+; stolonifera, 330+
BEER, S. C. *See* SHAW, J. & S. C. BEER, 112
BEER, S. C. *See* SHAW, J., S. C. BEER & J. LUTZ, 73
Bellemerea alpina, 106, 107+; cinereorufescens, 108; subsorediza, 68+
Bellia, 261
Biatora vernalis, 455
biosystematic study, of Sphagnum capillifolium, 1
BLACK, C. C. *See* GALLOWAY, C. M. & C. C. BLACK, 95
Blepharostoma trichophyllum, 350+, 500
BOWERS, F. D., R. R. HAYNES & S. L. TIMME. New or rare mosses of Alabama north of the Coastal Plain and Piedmont, 482
BOWERS, F. D., M. LELONG & A. R. DIAMOND. New and rare mosses of Alabama from the Piedmont and Gulf Coastal Plain, 485
Brachiolejeunea corticalis, *346*; insularis, *346*
Brachymenium systylium, 462
Brachythecium, 121, 159; calcareum, 474; curtum, 473; erythrorrhizon, 498; flexicaule, 474; leibergii, 194; mildeanum, 498; oedipodium, 473; plumosum, 498; salebrosum, 121+, 498; turgidum, 498
BRAKO, L. [Review of:] W. Korth and D. Ruckert, Bibliographie lichenologischer Bibliographien, 554
BRASSARD, G. R., M. FAVREAU & T. A. HEDDERSON. New or additional moss records from Newfoundland IX, 473
Braunia secunda, 463
Breedlovea chiapensis, 464
Breutelia, 373+; affinis, 369+*, 373; arcuata, 373; auriculata, 374+; chrysocoma, 369+*, 373; elongata, 369+*, 374+; grandis, 369+*, 374+; inclinata, 462; karsteniana, 369+*; pendula, 374; polygastrica, 369+*; robusta, 374; subarcuata, 462
British Columbia, Bryoerythrophyllum columbianum new to, 360; Mannia fragrans new to, 360; Pottia bryoides new to, 360; Pterygoneurum lamellatum new to, 360; Tortula caninervis new to, 360; Tortula scotteri new to, 360; Weissia brachycarpa new to, 361
Brodoa oroarctica, 108
Brotherella, 209+; complanata, 214; erythrocaulis, 214; falcata, 214; henonii, 214; lorentziana, 209+*; perpinnata, 214; propinqua, 214; pylaisiadelpha, 210; recurvans, 214; tenuirostris, 209
Bruchia brevifolia, new to Alabama, 130+, 260*; carolinae, 130; drummondii, 130; flexuosa, 130; fusca, 130; hallii, 130; ravenelii, 130, 486; texana, 130
Bryales, 271
Bryobrothera, 262+; crenulata, 264*
Bryocrumia andersonii, 183; vivicolor, 183

Bryoerythrophyllum campylocarpum, 463; columbianum, 360; inaequalifolium, 463; recurvirostrum, 463, 498
Bryohaplocladium microphyllum, 463; praelongum, 462
bryophytophagy, by crane fly larvae, 308+; of Rhizomnium punctatum, 308+
Bryopteris filicina, 330+; fruticulosa, 345
Bryoria, 82+, 108; chalybeiformis, 278, 501; furcellata, 275+, 278, 455; nitidula, 501
Bryum, 161; acutiforme, 498; argenteum, heavy metal tolerance in, 73, 74+, 113, 296; bicolor, 74, 178+, 309, 482, 486; billardieri, 463; caespiticium, 356, 498; capillare, 95+, 463, 498; cuspidatum, 309; lapponicum, 474; limbatum, 462; lisae var. cuspidatum, 74, 498; longisetum var. labradorense, 474; marratii, 473; ovatum, 473; petelottii, 463; procerum, 462; pseudotriquetrum, 463, 473, 498; purpurascens, 474; salinum, 474; schleicheri, 194; wrightii, 498
Buck, W. R. Henicodium replaces Leucodontopsis (Pterobryaceae), 534
Buck, W. R. [Review of:] M. Favreau & Guy Brassard, Catalogue Bibliographique des Bryophytes du Québec et du Labrador, 151
Buck, W. R. The status of the South American moss genus Syringothecium, 529
Buck, W. R. William Campbell Steere (1907–1989), 414
Buck, W. R. *See* Griffin III, D. & W. R. Buck, 368
Buck, W. R. & S. P. Churchill. The identity of Campylotrichum (Callicostaceae), 123
Buellia contermina, *68+*; disciformis, 275, 455; lacteoidea, 108+; papillosa, 107+; punctata, 275; spuria, 275; uberior, 68+
Buelliella, 68; trypethelii, 68
Bulbothrix isidiza, 68
Bullatina, 68; aspidota, 68+

Calamagrostis, 202
Calenia aspidotum, *68+*
Calicium corynellum, 68; trabinellum, 68
Callicosta bipinnata, 266
Callicostaceae, 123, 265+
Callicostella, 267
Calliergon, 128+, 161; cordifolium, 498; giganteum, 195, 498; richardsonii, 498; stramineum, 498; trifarium, 473, 475, 498
Calliergonella, 159
Calopadia, 68; fusca, 70; puiggarii, 70
Caloplaca ammiospila, 68+; bryochrysion, *68+*; caesiorufella, *68+*; castellana, 68+; cerina, 68, 275; c. fo. gilva, 68; cinnabarina, 275; cinnamomea, *68+*; citrina, 275; coralloides, 516; diphyodes, 68+; epiphyta, 68+; epithallina, 108+; feracissima, 275; flavorubescens, 275; gilva, *68*; holocarpa, 68+, 108, 275; invadens, *68+*; jungermanniae, 501; lithophila, *68+*; phaeocarpella, 68+; pyracea, *68+*; sideritis, 275; spitsbergensis, 68; stillicidiorum, *68+*, 108, 501; tetraspora, 108; tornoensis, 68
caloric content, of bryophyte sporophytes, 309
Calymperaceae, 302; further studies on Chinese, 170
Calymperes, 170+; afzelii, 170+, 303; cristatum, 170+; cucullatum, 170+; dozyanum, 170+; erosum, 170+; fasciculatum, 170+; fordii, 170+; graeffeanum, 170+; hampei, 170+; johanniswinkleri var. hasegawae, 170+; lonchophyllum, 170+; moluccense, 170+; serratum, 170+; strictifolium, 170+; tahitense, 170+; taitense, 170+; tenerum, 170+; tuberculosum, 170+
Calymperopsis, 170+; involuta, 170+; tjibodensis, 170+
Calypogeia caespitosa, 344; cellulosa, 330+; crenulata, 330+; cyclostipa, 344; elliottii, 330+; integristipula, 350+; laxa, 330+; lophocoleoides, 330+; peruviana, 330+; suecica, 350+
Calypogeja muelleriana, 500; sphagnicola, 500
Calyptopogon mnioides, 176
Calyptrochaeta, 262+; cristata, 266
Cameron, R. G. & R. Wyatt. Substrate restriction in entomophilous Splachnaceae: II. Effects of hydrogen ion concentration on establishment of gametophytes, 397
Campanula, 123; barbata, 123
Camptochaete, 159
Campylium chrysophyllum, 463, 474, 498; hispidulum, 498; h. var. sommerfeltii, 463; polygamum, 498; stellatum, 195, 498
Campylopodiella stenocarpa, 462
Campylopus bicolor, 299; introflexus, 301; lamprodictyon, 462; nivalis, 463; pilifer, 463; pittieri, 462; schimperi, 475; subturfaceus, 464
Campylostelium saxicola, new to Alabama, 482
Campylothelium nitidum, *69*
Campylotrichum, identity of, 123; acuminatum, *123*
Canada, Aloina bifrons new to, 359; changes to the lichen checklist of, 68; Crossidium aberrans new to, 359; Desmatodon convolutus new to, 359; Desmatodon guepinii new to, 359; Didymodon vinealis var. brachyphyllus new to, 359; Didymodon vinealis var. luridus new to, 361; Entosthodon rubiginosus new to, 359; soil conditions affecting community structure in, 495+; Tortula brevipes new to, 360; Trichostomopsis australasiae new to, 360
Candelaria concolor, 275; fibrosa, 275
Candelariella concolor var. effusa, 68; efflorescens, 275; vitellina, 108, 275
Cárdenas S., A. *See* Delgadillo M., C. & A. Cárdenas S., 461
Caribaeohypnum polypterum, 462
Catagonium, 159; brevicaudatum, 462
Catapyrenium, in North America, 190+; caeruleopulvinum, 192; cinereum, 192+; compactum, 190+; **daedaleum** sp. nov., 190+*; **globosum** sp. nov., 190+*; granulosum, 193; heppioides, 192; lachneum, 192; michelii, 192; plumbeum, 191+; **radiatum** sp. nov., 191+*; schaereri, 191*; squamellum, 193; tuckermanii, 192; waltheri, 192+*; zahlbruckneri, 192
Catillaria sphaeroides, 68+
Catoscopiaceae, 368+, *371*
Catoscopiées, *371*
Catoscopium, 371; nigritum, 369+*, 498
Caudalejeunea lehmanniana, 330+
Cephalozia affinis, 350+; bicuspidata, 500; caribbeania, *346*; connivens, 500; crassifolia, 330+; forficata, *346*; leucantha, 350+; lunulifolia, 500; pleniceps, 500; sandvicensis, *346*; subforficata, *346*
Cephaloziella antillana, 330+; arctica, 500; divaricata, 500; elachista, 500; hampeana, 500; rubella, 500; subdentata, 500
Cephaloziopsis intertexta, 330+
Ceratodon purpureus, 74, 112+, 356+, 498; stenocarpus, 463
Ceratolejeunea brevinervis, 330+; cornuta, 330+; cubensis, 346; flagelliformis, 330+; maritima, 330+; patentissima, 330+; portoricensis, *346*; rubiginosa, 330+; schwaneckei, *346*; spinosa, 330+; valida, 330+
Cetraria andrejevii, 501; aurescens, 275; ciliaris, 275; cucullata, 501; delisei, 501; ericetorum, 275+, 501; hepatizon, 107+, 501; islandica, 278, 501; laevigata, 501; nigricans, 501; nivalis, 501; platyphylla, 501; tilesii, 501

Chaenotheca gracilenta, 68+; phaeocephala, 68
Chaenothecopsis brevipes, 69
Chaetomitriopsis glaucocarpa, 266
Chaetomitrium, borbonicum, 267
Chamaecyparis nootkatensis, 118
Chandonanthus setiformis, 500
Cheilolejeunea adnata, 330+; clausa, 345; decidua, *346*; inflexa, 330+; involuta, 344; polyantha, 345; rigidula, 330+; trifaria, 330+
Chiapas, Aptychella new to, 461+; Grimmia new to, 461+
Chile, Orthotrichum truncatum sp. nov. from, 216
China, Bryocrumia andersonii new to, 183; Calymperaceae in, 170; comments on Horikawaea in, 490+; mosses new to, 183
Chionostomum, 529
CHRISTY, J. A. [Review of:] D. H. Vitt, J. E. Marsh & R. Bovey, Mosses, Lichens and Ferns of Northwest North America, 256
Chromatochlamys muscorum, 69
Chryso-hypnum diminutivum, 462
CHURCHILL, S. P. *See* BUCK, W. R. & S. P. CHURCHILL, 123
Cinclidium arcticum, 498; latifolium, 498; stygium, 195, 498; subrotundum, 498
Cirriphyllum cirrosum, 498
Cladidium bolanderi, 516
Cladina arbuscula, 275; rangiferina, 275
Cladonia, 82+, 126; subsect. Ochroleucae, 126; aberrans, 501; amaurocraea, 501; apodocarpa, 275; arbuscula, 501; bacillaris, 275, 501; bacilliformis, 501; bellidiflora, 501; caespiticia, 275; cariosa, 501; carneola, 126+, 501; cenotea, 501; chlorophaea, 275, 501; coccifera, 276, 501; coniocraea, 276, 501; cornuta, 276, 501; crispata, 126, 501; cristatella, 276, 312; cryptochlorophaea, 276; cyanipes, 501; cylindrica, 276; dactylota, 69; deformis, 501; ecmocyna, 501; fimbriata, 276, 501; furcata, 276; gracilis, 127, 278, 501; grayi, 276; homosekikaica, 69; humilis, 276; macilenta, 276; macrophylla, 501; maxima, 502; merochlorophaea, 502; mitis, 502; multiformis, 502; norrlinii, 502; parasitica, 276; peziziformis, 276; phyllophora, 108, 126, 502; piedmontensis, 127; pleurota, 502; polycarpoides, 69, 276; pyxidata, 108, 276, 502; ramulosa, 276; rangiferina, 502; rei, 276; robbinsii, 276; sp., 455; squamosa, 276, 502; stellaris, 502; subfurcata, 502; subulata, 278, 502; sulphurina, 502; uncialis, 502; verticillata, 276+, 502
Cladopodiella fluitans, 500
Claopodium, 159
Clathroporina diphloea, *69*+; exiguella, *69*+
Climacium, 309; dendroides, 95+, 498
Cnestrum alpestre, 474; glaucescens, 474
coevolution, bryophyte–invertebrate, 309
Collema nigrescens, 278; subflaccidum, 276; tenax, 276
Cololejeunea cardiocarpa, 330+; diaphana, *346*; minutissima, 330+; myriocarpa, *346*; obliqua, 330+; planifolia, 330+
Colura clavigera, 330+; rhynchophora, 330+; tortifolia, 330+
community structure, soil conditions affecting, 495+
Conocephalum conicum, 95+
Conostomoideae, 373+
Conostomum, 373+, 378; pentastichum, 369+*; tetragonum, 498
Conotrema urceolatum, 455
Cora pavonia, *69*+
Cornicularia aculeata, 502; divergens, 502; muricata, 502
Coscinodon calyptratus, 357; pilfera, *533*
Costa Rica, Sphagnum aureum in, 405+
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 131, 228, 420, 535
CRANDALL-STOTLER, B. & R. E. MAGILL. [Review of:] T. Koponen & D. H. Norris (eds.), Bryophyte Flora of the Huon Peninsula, Papua New Guinea, 151
Cratoneuron, 158; filicinum, 498
Crocynia gossypina, 69; neglecta, *69*
CROSBY, M. R. Recent literature on mosses—140, 245
Crosbya, 261+; straminea, 266
Crossidium aberrans, 359; asirense, 357; davidai, 357; rosei, 357+*
Crossomitrium, 265+; patrisiae, 266
Crossotolejeunea bermudiana, 330+; boryana, 330+
CRUM, H. New tropical American species of Sphagnum, 98
CRUM, H. & L. E. ANDERSON. New names for some North American mosses, 533
Cryphaea attenuata, 462; nervosa, 486; reticulata, 463
Cryptoleptodon flexuosus, 121
Ctenidium, 159; malacodes, 463, 474; molluscum, 474
Cyathodium steerei, 418
Cyathophoraceae, 261+
Cyathophorella, 262+; africa, 266
Cyathophorum, 262+
Cybebe gracilenta, *68*+
Cyclodictyon, 267; albicans, 264+*, 463
Cyclolejeunea accedens, 330+; angulistipa, *346*; chitonia, 330+; convexistipa, 330+; luteola, 330+; peruviana, 330+
Cynodon dactylon, 78+
Cynodontium strumiferum, 498; strumulosum, 474; tenellum, 498
Cyphelium trachylioides, 69
Cyrtolejeunea holostipa, 330+, 344
Cyrtomnium hymenophylloides, 498; hymenophyllum, 498
Cystolejeunea lineata, 330+

Dactylina arctica, 467+, 502; beringica, 467+; ramulosa, 502
Dactylospora parasitica, 69
Daltonia, 262+; gracilis, 269*, 462; lindigiana, 266, new to Mexico, 462; longifolia, 262; tenuifolia, new to Mexico, 462
Daltoniaceae, 261+
DEGUCHI, H. *See* LEWINSKY, J. & H. DEGUCHI, 216
DELGADILLO, M., C. Astomiopsis × altivallis (Musci: Ditrichaceae), a putative interspecific hybrid in Mexico, 225
DELGADILLO M., C. & A. CÁRDENAS S. Phytogeography of high-elevation mosses from Chiapas, Mexico, 461
Dendroceros crispus, 331+
Dendrocyathophorum, 271
Dendrographa minor, 516
Dendropogonella rufescens, 462
Dermatocarpon daedaleum, *190*; luridum, 276; miniatum, 191, 276; monstrosum, *191*+; waltheri, *192*
Desmatodon convolutus, 357; guepinii, 359; heimii var. arctica, 498; latifolius, 359; leucostoma, 359; obtusifolius, 359; plinthobius, 486; steereanus, 418; suberectus, 359
DIAMOND, A. R. *See* BOWERS, F. D., M. LELONG & A. R. DIAMOND, 485
Dianthelia steerei, 418
diaspores, production of, 351+; sizes of, 351+
Dicranella grevilleana, 498; hookeri, 177; schreberiana var. robusta, 475; sharpii, 462
Dicranum, 121; acutifolium, 195, 498; amannii, 498; angustum, 498; brevifolium, 498; condensatum, 314+;

elongatum, 498; flagellare, 463; frigidum, 462; fuscescens, 498; groenlandicum, 498; majus, 498; muehlenbeckii, 498; rhabdocarpum, 314+; scoparium, 121, 314+, 498; spadiceum, 498; undulatum, 498
Dictyonema glabratum, 69+; pavonium, *69+*
Didymodon fallax, 463; reedii, 359; rigidulus, 463, 498; tophaceus, new to Alabama, 486; vinealis complex, 359; v. var. brachyphyllus, 359; v. var. luridus, 359+
Dimelaena oreina, 106+; radiata, 521
Dimorphocladon borneense, 267
Diphyscium foliosum, 309, 324+
Diplasiolejeunea brachyclada, *346*; cavifolia, 330+; pellucida, 330+; rudolphiana, 330+; unidentata, 330+
Diploneuron, 270
Diploschistes actinostomus, 69; albissimus, *69*; bryophilus, *69*; canadensis, *69+*; diacapsis, 69+; gypsaceus, 69; hypoleucus, 69+; muscorum, 69; scruposus, 108; straminescens, *69+*
Distichium capillaceum, 498; inclinatum, 498
Distichophyllaceae, 261+
Distichophyllidium, 262+
Distichophyllum, 261+, 262+; crispulum, 266+; freycinetti, 264*; pulchellum, 266; rotundifolium, 266
Ditrichum crispatissimum, 475; flexicaule, 475, 498; giganteum, 475; lineare, 314+; pallidum, 314+; pusillum, 486
Dominican Republic, Drepanocladus vernicosus in, 128
Donnellia, *529*
Drepanocladus, 128+; aduncus, 196, 498; crassicostatus, *533*; exannulatus, 195, 498; e. var. mexicanus, 462; fluitans, 498; intermedius, 161; lapponicus, 161; lycopodioides var. brevifolius, 498; pseudostramineus, 498; revolvens, 161, 195, 498; sendtneri, 196+; uncinatus, 499; vernicosus, 128+, 161, 499
Drepanolejeunea anoplantha, 330+; biocellata, 330+; crassiretis, 330+; crucianella, 330+; dissitifolia, 330+; evansii, 330+; fragilis, 330+; inchoata, 330+; infundibulata, 346; lichenicola, 330+; mosenii, 330+; orthophylla, 330+; pinnatiloba, 330+; trigonophylla, 330+
Drimys winteri, 176
Dumortiera hirsuta, 331+; nepalensis, *346*
DURING, H. J. *See* WERGER, M. J. A. & H. J. DURING, 411
Duthiella, 159

Echinodium, 159
Ectropothecium, 210
EGAN, R. S. Changes to the "fifth checklist of the lichen-forming, lichenicolous and allied fungi of the continental United States and Canada." Edition I, 68
EGAN, R. S. Recent Literature on Lichens—135, 141;—136, 239;—137, 427
EGAN, R. S. [Review of:] L. Tibell, Australasian Caliciales, 258
eicosapentaenoic acid, in lipids of Bryum bicolor, 178+
electrophoresis, procedure for comparative analysis, 164
ELLIOTT, J. C. & G. L. MOORE. Additions to the moss flora of Montana, 194
EL-SAADAWI, W. E. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 178
Encalypta affinis, 499; a. var. **macounii** comb. nov., 533; alpina, 475+, 499; brevicolla subsp. crummiana, *533*; b. var. **crummiana** comb. nov., 533; macounii, *533*; procera, 473, 499; rhaptocarpa, 499; streptocarpa, 473
Endocarpon daedaleum, *190*; pusillum, 276
Entodon, 122, 529; beyrichii, 463; brevirostris, 463; jamesonii var. mexicanus, 462; seductrix, 95
Entodontopsis pygmea, 183
Entosthodon drummondii, 486; obtusifolius, 463; rubiginosus, 359; steerei, 418
enzymes, of sucrose metabolism, 95
Eopyrenula leucoplaca, 276
Ephemeropsaceae, 261+
Ephemeropsis, 267+; tjibodensis, 267
Ephemerum crassinervium, 486; spinulosum, new to Alabama, 486
epiphytes, succession on Quercus garryana, 81
epixylic bryophytes, habitat requirements of, 349+
Eriopus, 262
Erythrodontium longisetum, 463
Eubryales, 270+
Eucladium verticillatum, 196+; new to Alabama, 487
Euosmolejeunea clausa, *346*; duriuscula, *346*; rigidula, *346*; trifaria, *346*
Eurhynchium pulchellum, 463, 499; riparioides, 463, 482
Europe, Pseudoleskeella in, 387+
Evernia divaricata, 502; mesomorpha, 276, 455; perfragilis, 502; prunastri, 82+
evolution, of heavy metal tolerance, 73; peristome development in relation to, 314+
EWAN, J. [Review of:] R. W. Sims, P. Freeman & D. L. Hawksworth, Key Works of the Fauna and Flora of the British Isles and North-western Europe, 554
Exodictyon, 493

FAVREAU, M. *See* BRASSARD, G. R., M. FAVREAU & T. A. HEDDERSON, 473
Fissidens sect. Aloma, 523+; sect. Fissidens, 523+; sect. Pachylomidium, 523+; sect. Semilimbidium, 523+; subgen. Fissidens, 523+; subsect. Bryolimbidium, 523+; adianthoides, 499; asplenioides, 463, 525; asterodontius, 523; austro-americanus, 524; brasiliensis, *525+*; flabellatus, 525; flexinervis, 523; fontanus, 196, 463; grandifrons, 196; guianensis, 526; hornschuchii, 525+; intermedius, 526; legalloi, *526*; leptophyllus, 523+; fo. obtusissimus, *524*; martianus, *526*; **obtusissimus** stat. nov., 524; osmundoides, 499; repandus, 463; serrulatus, *525+*; sharpii, 526; steerei, 418; virens, 183
Fistulariella, 69; inflata, 71
FLATBERG, K. I. Sphagnum (Cuspidata) pacificum, sp. nov., 116
Flavoparmelia baltimorensis, 326+; caperata, 455
Fleischerobryum, 374; longicolle, 369+*
Floribundaria floribunda, 121+
Florschuetziella steerei, 418
Flowersia gen. nov., 372+; **abyssinica** comb. nov., 372+; **campylopus**, 369+*, comb. nov., 372+; **setifolia**, 369+*, comb. nov., 372+; **sinensis**, 369+*, comb. nov., 372+
Folioceros, 219+; assamicus, 219+; dixitianus, 219+; glandulosus, 219+; indicus, 219+; **kashyapii** sp. nov., 219+*; mangaloreus, 219+; pandei, 219+; physocladus, 219+; pinnilobus, 219+; satpurensis, 219+; spinisporus, 219+; udarii, 219+*
Fontinalis antipyretica, 308; hypnoides, 196; sullivantii, new to Alabama, 483; welchiana, new to Alabama, 483
Foreauella orthothecia, 183
FORERO, E. [Review of:] J. L. Hartwell. Plants Used Against Cancer. A Survey, 153
Fossombronia brasiliensis, 331+
FRANCIS, P. C. & R. L. PETERSEN. Assessment of toxicity of heavy metal ion combinations on spore germination and protonemal growth of Polytrichum commune, 60
Frullania, in Western North America, 363+; sect. Frullania, 363+; sect. Integristipulae, 365; sect. Mayebarae, 365; sect. Trachycolea, 366+; subgen. Frullania,

363+; subgen. Trachycolea, 363+; apiculata, 330+; arecae, 330+; arietina, *346*; atrata, 330+; bolanderi, 83+, 363+; brasiliensis, 330+*; brittoniae, 363+; californica, 363+; catalinae, 365+; chilcootiensis, 363+; eboracensis, 363+; ericoides, 330+; exilis, 345; franciscana, 363+; gibbosa, 330+; hattoriana, 363; hians, *346*; inflata, 363+; jackii, 365; kunzei, 330+; major, 363; mayebarae, 365; nisquallensis, 363+; nodulosa, 330+; obcordata, 330+, *346*; riojaneirensis, 330+; riparia, 363+; squarrosa, *346*; steereana, 418; subtilissima, 345; tamarisci, *363*
Frullanoides corticalis, 330+; liebmanniana, 330+; tristis, *346*
Fulgensia bracteata, 502
Funaria, 64+; hygrometrica, 60+, 74, 95+, 114, 309; var. calvescens, 463
Fuscidea placidensis, *69*

GALLOWAY, C. M. & C. C. BLACK. Enzymes of sucrose metabolism in bryophytes, 95
Gammiella, 214
Geocalycaceae, 123
GIANCOTTI, C. & D. M. VITAL. Lejeunea capensis Gott. (Hepaticae: Lejeuneaceae) disjunct between South America and Africa, 305
GLIME, J. M. [Review of:] T. H. Nash, III & V. Wirth (eds.), Lichens, Bryophytes and Air Quality, 448
Gloeocapsa, 25
Glossadelphus bilobatus, 184; glossoides, 184; laevifolius, 184
Glyphocarpa abyssinica, *372*; cernua, 374, *376*; glaziovii, *376*; lindigii, *376*; pusilla, *376*
Glyphocarpus setifolius, *372*
Gollania, 160
GOWAN, S. P. The lichen genus Porpidia (Porpidiaceae) in North America, 25
GRADSTEIN, S. R. A key to the Hepaticae and Anthocerotae of Puerto Rico and the Virgin Islands, 329
GRANZOW-DE LA CERDA, I. Notes on five species of Anomodon, some with erroneous identity, including two new combinations, 381
Graphis scripta, 276, 455
GRIFFIN III, D. & W. R. BUCK. Taxonomic and phylogenetic studies on the Bartramiaceae, 368
Grimmia affinis, 475+; alpestris, 357; anomala, 473; apocarpa fo. latifolia, *533*; atrata, 475; donniana, 473; hartmannii var. anomala, 473; incurva, 195; laevigata, 196, 357; montana, 357; ovalis, 357, 475; trichophylla, 463
Groutiella chimborazense, 462; tomentosa, 184
GUNTHER, A. J. Nitrogen fixation by lichens in a subarctic Alaskan watershed, 202
Gyalectidium rotuliforme, *68+*
Gyalidea lecideopsis var. convarians, 69
Gyalideopsis alnicola, 69
Gymnocolea inflata, 165, 500
Gymnomitrion concinnatum, 500
Gymnostomum aeruginosum, 463, 499; setifolium, *372*

Haematomma lapponicum, *69+*, 106+; pustulatum, 455; rappii, *69+*; ventosum, *69+*
Halecania, 69; alpivaga, 69+
HAMADA, N. The effect of various culture conditions on depside production by an isolated lichen mycobiont, 310
HAMMER, S. Cladonia carneola: Two new localities in western North America, 126
Haplohymenium, 381; triste 383
haplolepideae, peristome development in, 314+
Harpalejeunea heterodonta, *346*; subacuta, 330+; tridens, 330+; uncinata, 330+
Harpidium exannulatum var. falcatum, *533*
HAYNES, R. R. *See* BOWERS, F. D., R. R. HAYNES & S. L. TIMME, 482
HE, S. *See* REDFEARN, JR., P. L., P.-C. WU, S. HE & Y.-G. SU, 183
heavy metals, 60+; tolerance in a moss, 73
HEDDERSON, T. A. *See* BRASSARD, G. R., M. FAVREAU & T. A. HEDDERSON, 473
HEDENÄS, L. Drepanocladus vernicosus in the Dominican Republic, 128
HEDENÄS, L. Some neglected character distribution patterns among the pleurocarpous mosses, 157
Hedwigia, 265; ciliata, 463
Hedwigidium integrifolium, 464
Helicoblepharum, 268+
Helodium paludosum, new to Alabama, 487
Hemiragis, 270; aurea, 266
Henicodium, 534+; **geniculatum** comb. nov., 534+; niamniamiae, *534*
Hepaticae, chemosystematics of, 164; of Puerto Rico and the Virgin Islands, 329+
Herberta juniperina, *346*
Herbertus, 121; divergens; 329+; elliottii, 344; juniperoideus, 329+; pensilis, 329+
herbivore, of lichens, 326+; of moss, 308
Herpetineuron acutifolium, 385+*; toccoae, 385, 464, new to Alabama, 483
Herzogiella, 159; cylindricarpa, 462
Heterocladium, 159, 387, 529
Heterodermia granulifera, 276; leucomelos, 278; obscurata, 276; sp., 516; speciosa, 276
Heterophyllium affine, 464
Heteroscyphus elliottii, 330+
Himalaya, bryophytes used in northwest, 120
Holomitrium flexuosum, 462; leptopodum, 286; longipes, 286+; perichaetiale, 286
Homaliadelphus sharpii var. sharpii, 462
Homalothecium fulgescens, 82+; leskeoides, 463; nuttallii, 82+
HONG, W. S. The genus Frullania in North America west of the hundredth meridian, 363
Hookeria, 261+; acutifolia, 264*; group, 267+; lucens, 266+, 269*
Hookeriaceae, 261+
Hookeriales, 262+; familial classification of, 261
Hookeriopsis, 265; parkeriana, 269*; steerei, 418; undulata, 266
Horikawaea dubia, 490+*; nitida, 490+*
Horridohypnum mexicanum, 462
HORTON, D. G. [Review of:] J. M. Glime (ed.), Methods in Bryology, 255
Hubbsia californica, 69; lumbricoides, *69*
Huilia, 25+; phylliscina, 57; platycarpoides, *69+*; sorediozodes, 58
hybrid, interspecific moss, in Mexico, 225
Hygrohypnum steerei, 418
Hygrolejeunea cerina, *346*; orba, 346
Hylocomium, 120; splendens, 499
Hymenelia lacustris, 276
Hyocomium, 159
Hyophila involuta, 464
Hyperphyscia adglutinata, 276; syncolla, 276
Hypnaceae, 529
Hypnales, 270+
Hypnella, 270
Hypnodendron, 198+; auricomum subsp. auricomum, 198+*; a. subsp. opacum, *198+*; a. subsp. sumatra-

num, 198+*; dendroides, 198; diversifolium, 198; milnei subsp. korthalsii, 198; opacum, a reevaluation of, 198+; reinwardtii subsp. caducifolium, 198; subspininervium subsp. arborescens, 198; sumatranum, *200*; vitiense, 198
Hypnum, 120, 159, 211+; sect. Henicodium, 534; bambergeri, 499; callichroum, 475; cupressiforme, 121+, 308, 499; c. var. filiforme, 210, 475; c. var. lacunosum, 475; curvifolium, 462; fluitans var. falcatum, *533*; hamulosum, 499; holdridgei, 210; jutlandicum, 475; lindbergii, 499; mammillatum, 473; niam-niamiae, *534*; pylaiseaceum, nom. nud., 210; revolutum, 499; tectorum, *391*; uncinatum subsp. symmetricum, *533*; vaucheri, 499; viridicolor, nom. nud., 210
Hypocenomyce scalaris, 276
Hypogymnia austerodes, 108, 502; enteromorpha, 83+; heterophylla, 82+; physodes, 82+, 455, 502; subobscura, 502; tubulosa, 82+
Hypopterygiaceae, 271+
Hypopterygium, 271; commutatum, 271; faurei, 271; tamariscinum, 463
Hypotrachyna pseudosinuosa, 69; sinuosa, 83+

Icmadophila ericetorum, 502
Imshaugia aleurites, 276
India, two species of Folioceros from, 219
Inflatolejeunea capensis, *306*
Interspecific hybrid, in Mexico, 225
Isobryales, 271
Isopterygium, 159, 529+; **brasiliense** comb. nov., 531+*; distichaceum, 475, new to Alabama, 483; elegans, 475; jamaicense, 529; pulchellum, 499; **sprucei** comb. nov., 529+*
Isotachis erythrorhiza, 330+; multiceps, 330+; serrulata, 330+
Isothecium, 121, 159; alopecuroides, 473; stoloniferum, 83+

Jamesoniella colorata, *346*; rubricaulis, 330+
Jasper National Park, saxicolous lichens of, 105
JOHN, E. A. The saxicolous lichen flora of Jonas Rockslide, Jasper National Park, Alberta, 105
Julella, 71; sublactea, 69
Jungermannia, 123; callithrix, 330+; cerina, *345*; dominicensis, *346*; leiantha, 165

KERR, S. & M. S. ZAVADA. The effect of the lichen Acarospora sinopica on the elemental composition of three sedimentary rock substrates in South Africa, 407
Kiaeria blyttii, 475; glacialis, 499
Kirschsteiniothelia aethiops, 275
Kurzia capillaris, 330+; verrucosa, *346*
KUSHNER, B. K. *See* BASILE, D. V., B. K. KUSHNER & M. R. BASILE, 164
Kymatocalyx dominicensis, 330+

Lasallia papulosa, 326+; pensylvanica, 108
Laurera megasperma, 69+
LAWREY, J. D. Lichen secondary compounds: Evidence for a correspondence between antiherbivore and antimicrobial function, 326
Lecania alpivaga, *69*+; arctica, *68*+; thallophila, *69*+
Lecanora subgen. Placodium sect. Endochloris, 513+; allophana, 276; argopholis, 69, 109; bolanderi, 513; californica, 516; campestris, 276; cenisea, 108; demissa, 513; deustata, 70; diffracta, *69*; dispersa, 276; epibryon, 502; frustulosa, *69*, 108; hagenii, 276; intricata, 108; intrudens, *69*+; lisbonensis, 513; marginata, 69+; muralis, 69+, 276; occidentalis, *69*; oregana, *69*; **pachysoma** sp. nov., 518+*; pallida, 276, 455; phryganitis, 515+*; pinguis, 516+*; polytropa, 108, 276; rupicola, 108; strobilina, 455; thysanophora, 455; varia, 276, 455; **xanthosora** sp. nov., 520+*
Lecidea, 25+, 69; aenea, *69*+; amylacea, *69*+; atrobrunnea, 108+; auriculata, 108; cinereoatra, *39*; circumnigrata var. reagens, *69*+; confluens, 108; contraponenda, *41*; crustulata, *41*; cyrtidia, 276; deustata, *69*; diversa, *42*+; elata, *69*+; erratica, 276; flavicunda, *43*+; flavocoereulescens, *46*+; furvella, *69*+; garovaglii, *69*+; glaucophaea, *47*; impavida, *69*+; instrata, *69*+; insularis, 71; intumescens, 71; lactea, 276; lapicida, 108; fo. ochracea, 108; leucophaea, *69*+; lithophila, 108; lulensis, *69*+; marginata, *69*+; pallida, *69*+; parasema var. crustulata, *41*; paupercula, 106, 108+; placidensis, *69*; plana, 108+; purissima, *69*+; rubiformis, 502; soredifera, *41*+; subplumbea, *69*+; subsorediza, *68*+; superba, *53*; tessellata, 108; vernalis, 276; zeoroides, *56*
Lecidella carpathica, 108; placidensis, 69+; stigmatea, 107+
Leciographa inspersa, 69; parasitica, *69*+
Leioderma sorediatum, 69
Leiomela, 371+; bartramioides, 369+*; setifolia, *372*
Leiosporoceros, 219+
Lejeunea bethanica, 346; capensis, 305+*; cerina, *346*; cladogyna, 330+; elliottii, 346; flava, 330+*; floridana, 330+; glaucescens, 330+; laetevirens, 330+; minutiloba, 330+; pililoba, *346*; trinitensis, 330+
LELONG, M. *See* BOWERS, F. D., M. LELONG & A. R. DIAMOND, 485
Lembophyllum, 159
Lepidolejeunea involuta, 330+
Lepidopilum, 268+; polytrichoides, 123; pringlei, 462; tortifolium, 266
Lepidozia capillaris, *346*; commutata, *346*; patens, 330+; portoricensis, 344; reptans, 350+
Lepraria finkii, 276; membranacea, 276; neglecta, 69+, 107+, 276; sp., 455, 516
Leprocaulon subalbicans, 107+
Leptobryum pyriforme, 499
Leptocolea cardiocarpa, *346*; planifolia, *346*; scabrifolia, *346*
Leptodictyum **humile** comb. nov., 533; kochii, 533; trichopodium, 533
Leptodontium flexifolium, 464; viticulosoides, 464; var. sulphureum, 463
Leptogium, 82+; burnetiae, 276; corticola, 276; cyanescens, 276; dactylinum, 276; gelatinosum, 82+; juniperinum, 276; lichenoides, 203, 276, 502; saturninum, 502
Leptolejeunea elliptica, 330+; exocellata, 330+; hamulata, *346*; stenophylla, *346*
Leptoscyphus gibbosus, 330+
Leptotrema reclusum, *70*+
Lescuraea, 162; **arizonae** comb. nov., 395+; incurvata, *388*; saxicola, 475; stenophylla, 194
Leskea gracilescens, 487; nervosa var. sibirica, *390*; papillosa, *395*; polycarpa, 475; tenuirostris, *210*
Leskeella, 387+; arizonae, *395*; nervosa var. sibirica, *390*
Leskeodon, 262+; pusillus, 266
Leskeodontopsis, 262+
Leucobryum glaucum, 493+; martianum, 493+*; polakowskyi, 463
Leucodon curvirostris, 462; geniculatus, *534*; sciuroides, 121
Leucodontaceae, 534
Leucodontales, 270
Leucodontopsis, *534*; geniculata, *534*; niam-niamiae, *534*

Leucolejeunea unciloba, 330+; xanthocarpa, 330+
Leucomium, 267, 529; steerei, 419; strumosum, 264*
Leucophanes, 493
Leucosarmentum portoricense, *346*
LEWINSKY, J. & H. DEGUCHI. Orthotrichum truncatum Lewinsky & Deguchi, sp. nov. from Chile, and a new record for O. macrosporum Lewinsky, 216
Lichen albo-caerulescens, *35*; speireus, *51*
lichen mycobiont, depside production by, 310+
Lichenothelia, 70; calcarea, 70; convexa, 70; intermixta, 70; metzleri, 70; scopularia, 70; tenuissima, 70
LIN, P.-J. & W. D. REESE. A further study on Chinese Calymperaceae, 170
LIN, P.-J. *See* REESE, W. D. & P.-J. LIN, 186
Lindbergia mexicana, 462
lipids, of Bryum bicolor, 178+
Lobaria oregana, 205; pulmonaria, 83+, 203+, 276; quercizans, 276; scrobiculata, 83+
Loeskypnum badium, 499; wickesiae, 475
Lopadiopsis floridana, *68*+
Lopadium fuscum, *70*; puiggarii, *70*
Lophocolea bidentata, 330+; coadunata, *346*; connata, 330+; cubana, *346*; guadeloupensis, 330+; heterophylla, 350+; martiana, 330+; minor, 500; sintenisii, *346*; trapezoidea, 330+
Lopholejeunea howei, *347*; muelleriana, 330+; quelchii, 330+; sagraeana, *347*; subfusca, 330+
Lophozia alpestris, 500; ascendens, 350+, 500; badensis, 500; collaris, 500; excisa, 500; gillmanii, 500; grandiretis, 500; guttulata, 500; heterocolpos, 500; heteromorpha, 500; incisa, 350+; longidens, 350+, 500; longiflora, 350+; obtusa, 500; opacifolia, 501; rubrigemma, 501; rutheana, 501; schistophila, *347*; silvicola, 350+; sudetica, 501; ventricosa, 501; wenzelii, 501
Luisierella barbula, new to Alabama, 487
Lunularia cruciata, 164
LUO, J.-X. Comments on the moss genus Horikawaea (Pterobryaceae) in China, 490
LUTZ, J. *See* SHAW, J., S. C. BEER & J. LUTZ, 73

Macrocoma, 296; tenue subsp. sullivantii, 464
Macrolejeunea **cerina** comb. nov., 345+; subsimplex, *345*, *347*
Macromitrium, 217; brownii, *290*; caducipilum, 286; cirrosum, 282; fragilicuspis, 462; guatemalense, 462; hectorii, *282*; longifolium, 463; longirostre, 286; moorcroftii, 121
Macrothamnium submacrocarpum, 121+
MAGILL, R. E. *See* ALLEN, B. H. & R. E. MAGILL, 543
MAGILL, R. E. *See* CRANDALL-STOTLER, B. & R. E. MAGILL, 451
MAGILL, R. E. & B. H. ALLEN. Recent literature on mosses—141, 434
Mannia fragrans, 360; pilosa, 360
Marchantia chenopoda, 331+; domingensis, *347*; inflexa, 331+; palmata, 120; polymorpha, 60+, 74+, 120, 501
Marchesinia brachiata, 330+
Maronea porinoidea, *70*+
Masonhalea richardsonii, 502
Mastigolejeunea auriculata, 330+
MATTERI, C. M. Additions to the moss flora of Tierra del Fuego (Argentina), 174
MCINTOSH, T. T. Bryophyte records from the semiarid steppe of northwestern North America, including four species new to North America, 356
MCQUEEN, C. B. A biosystematic study of Sphagnum capillifolium sensu lato, 1
MCQUEEN, C. B. Sphagnum aureum, a new species from Costa Rica, 405
Meesia triquetra, 195+, 369+*, 499; uliginosa, 369+*, 499
Meesiaceae, 368+
Megaceros alatifrons, 344; vincentianus, 331+
Meiotheciopsis, 529
Melanelia granulosa, 108; panniformis, 108; sorediata, 108; stygia, 108+; subaurifera, 82+, 455
Mesoptychia sahlbergii, 501
metabolism, of sucrose in bryophytes, 95
Meteoriaceae, 268
Meteoridium remotifolium, 463
Meteoriopsis reclinata, 121
Meteorium illecebrum, 463
Metzgeria albinea, 344; australis, 344; decipiens, 331+; dichotoma, 344; furcata, 344; hamata, *347*; leptoneura, 331+*; oligotricha, *347*; procera, 331+; sandei, 344; uncigera, 331+; vivipara, 331+
Mexico, Daltonia lindigiana new to, 461+; Daltonia tenuifolia new to, 461+; interspecific hybrid in, 225; Orthotrichum trachymitrium new to, 461+; phytogeography of high-elevations in, 461+
Micarea melaena, 276
Microglaena, 70; hassei, *70*+; inductula, *70*+; muscorum, *69*; sordidula, *70*+; subcorallina, *70*+
Microlejeunea acutifolia, 330+; bullata, 330+; laetevirens, *347*; monoica, 330+
Microlepidozia capillaris, *347*
Micromitrium austinii, new to Alabama, 487; megalosporum, new to Alabama, 487
Micropterygium carinatum, 330+; exalatum, 344; portoricense, *347*
Microthelia aterrima, *70*; metzleri, *70*; sychnogonoides, *70*+
Miriquidica, 70; deusta, 69+; garovaglii, 69+; griseoatra, 69+; instrata, 69+; intrudens, 69+; leucophaea, 69+; lulensis, 69+
MISHLER, B. D. *See* SHAW, J., B. D. MISHLER & L. E. ANDERSON, 314
Mittenia plumula, 314
Mittenothamnium reptans, 463
Mitthyridium, 170+; fasciculatum, 170+; flavum, 170+
Mniaceae, 308
Mnium affine, 95+; a. var. rugicum, 480; blyttii, 499; medium, 480; punctatum, 479; p. subsp. chlorophyllosum, 479; p. var. elatum, 479; rugicum, 195, 480; thomsonii, 499
Molendoa sendtneriana, 464, 473
Monoclea gottschei, 331+
Montana, additions to moss flora of, 194
MOORE, G. L. *See* ELLIOTT, J. C. & G. L. MOORE, 194
Muelleriella crassifolia, 296
MURRAY, B. M. [Review of:] V. D. Aleksandrova, Vegetation of the Soviet Polar Deserts, 451
Muscites steerei, 419
Mycocalicium subtile, 276
Myelochroa, 70; aurulenta, 70+; galbina, 70+, 455; metarevoluta, 70; obsessa, 70
Mylia anomola, 501; gibbosa, *347*
Myriotrema reclusum, 70+
Myurella, 160; julacea, 499; tenerrima, 476, 499

Nardia callithrix, *347*; succulenta, 330+
Neckera chlorocaulis, 462; crenulata, 121; douglasii, 83+; ehrenbergii, 462; menziesii, 82+
Neckeropsis crinita, 184
Neesioscyphus bicuspidatus, 330+
Neomeesia paludella, 369+*
Nephroma, 203+; arcticum, 203, 502; bellum, 203, 278; expallidum, 502; helveticum, 502; resupinatum, 82+

Neurolejeunea breutelii, 330+; catenulata, 330+; **sastreana** sp. nov., 345+*
Newfoundland, Pohlia atropurpurea, new to, 476; Pseudoleskeella papillosa new to, 476; additions to the moss flora of, 473+
New York, Rhizomnium pseudopunctatum in, 479+
New Zealand, the genus Schlotheimia in, 282+
Niebla sp., 519
nitrogen fixation, by lichens in Alaska, 202
Normandina pulchella, 82+
NORRIS, D. H. *See* WILSON, P. & D. H. NORRIS, 387
North America, additions and revised key to Catapyrenium in, 190; bryophytes new to, 356; changes to lichen checklist of, 68; Crossidium rosei new to, 357; Frullania in, 363+; Phascum vlassovii new to, 359; Pottia wilsonii new to, 357; Pseudoleskeela in, 387+; Pterygoneurum kozlovii new to, 357; Porpidia in, 25; new localities of Cladonia carneola in western, 126
Notothylas breutelii, 331+
Nowellia bicornis, *347*; caribbeania, *347*; dominicensis, 330+; evansii, 330+

Ochrolechia androgyna, 108, 502; frigida, 502; pallescens, 276; sp., 516; upsaliensis, 502
Octoblepharum, 493
Odontolejeunea lunulata, 330+; sieberiana, *347*
Odontoschisma denudatum, 344; longiflorum, 344; macounii, 501; portoricense, *347*; prostratum, 330+
Ohio, Cuyahoga Valley National Recreation Area of, 273+; lichens and air quality in, 273
Oligotrichum aligerum, 195; hercynicum, 476
Omphalanthus filiformis, 330+
Oncophorus virens, 499; wahlenbergii, 499
Onoclea sensibilis, 60+
Opegrapha varia, 276; viridis, 277; vulgata, 277
Ophioparma, 70; lapponica, 69+, 70; ventosa, 70
Oregon, Didymodon vinealis var. luridus new to, 361; epiphyte succession in the Willamette Valley of, 81
Orphniospora moriopsis, 108
Orthothecium chryseum, 499; intricatum, 499; strictum, 499; chryseum, 476
Orthotrichum, 216+; sect. Rupestria, 218; subg. Phaneroporum, 218; aequatoreum, 462; angulosa, 282; assimile, 216+; bicolor, 216; brotheri, 216; densum, 218+; elegantulum, 216+; hortense, 216+; laevigatum, 357; ludificans, 216+; lyellii, 83+; macrosporum, 216+; ohioense, 487; pallens, 476; pallidum, 282; perexiguum, 216; pycnophyllum, 462; rupestre, 216, 473; steerei, 419; stramineum, 473; trachymitrium, new to Mexico, 462; tristriatum, 216; **truncatum** sp. nov., 216+*
Osmunda cinnamomea, 60+; claytoniana, 60+
Oxystegus **spiralis** comb. nov., 533; tenuirostris, 464, 499

Pachyspora verrucosa, 502
Palaeohypnum steerei, 419
Pallavicinia lyellii, 331+
Paludella squarrosa, 369+*
Pannaria, 82+; leucophaea, 70; microphylla, *70*
Papillaria deppei, 463; fuscescens, 121; nigrescens, 464
Paracromastigum bifidum, 330+
Parmelia antillensis, *70*; aurulenta, *70*+, 277; bolliana, 277; caperata, 277; congensis, *70*+; coniocarpa, 277; conspersa, 278; crinita, 278; dissecta, *70*; fertilis, 70; finkii, *70*+; flaventior, 277; galbina, *70*+, 277; horrescens, *70*; hubrichtii, *70*+; kerguelensis, 70; livida, 277; margaritata, 277; metarevoluta, *70*+; neodiscordans, 70; obsessa, *70*+; olivacea, 278; omphalodes, 70, 108, 502; o. subsp. glacialis, *70*; o. subsp. omphalodes, 70; perforata, 278; plittii, 277; pseudosulcata, *70*; rudecta, 274+; saxatilis, 108, 502; schaereri, *191*; silvestris, *70*+; skultii, 70+; spumosa, *70*+; squarrosa, 277, 455; stygia, 502; subaurifera, 277; subquercifolia, *70*+; subrudecta, 277; subtinctoria, 277; sulcata, 82+, 277, 455
Parmeliella microphylla, *70*
Parmelina antillensis, *70*+; aurulenta, *70*+; dissecta, *70*+; galbina, *70*+; horrescens, *70*+; metarevoluta, *70*+; minarum, *70*+; obsessa, *70*+; spumosa, *70*+
Parmelinopsis, 70; horrescens, 70; minarum, 70+; spumosa, 70+
Parmeliopsis ambigua, 455
Parmotremopsis, 70; antillensis, 70+
Patellaria macrocarpa, *50*
Pellia endiviifolia, 501; epiphylla, 308
Peltigera, 203+; aphthosa, 203, 278, 502; canina, 277, 502; collina, 82+, 203; didactyla, 277; evansiana, 277+; horizontalis, 277, 502; malacea, 203, 502; polydactyla, 203, 277, 502; rufescens, 278; scabrosa, 203
Pennsylvania, Scopelophila cataractae in, 112
Penzigiella cordata, 184
peristome, Bryum type, 315+; daltoniaceous type, 267+; development of, 314+; diplolepideous type, 314+; Funaria type, 315+; haplolepideous type, 314+; hookeriaceous type, 267+; hygrocastique type, 268+; hygroscopic motion of, 268; xerocastique type, 268+
Pertusaria dactylina, 502; leucostoma, 277; macounii, 455; opthalmiza, 277; pertusa, 278; pustulata, 277; trachythallina, 455; velata, 277
PETERSEN, R. L. *See* FRANCIS, P. C. & R. L. PETERSEN, 60
Phaeoceros laevis, 331+
Phaeophyscia adiastola, 277; ciliata, 277; endococcinea, 108; orbicularis, 277; pusilloides, 277; rubropulchra, 277, 455; sciastra, 108
Phascum cuspidatum, 357; smardaeanum, 357; vlassovii, 359+*
phenolic compounds, 326+
Philonotis, 372+; sect. Catenularia, 375; sect. Leiocarpus, 375+; sect. Philonotula, 176; **australiensis** nom. nov., 376; **bartramioides** comb. nov., 376; **cernua** comb. nov., 376; **comosa** comb. nov., 376; corticata, 375; **curta** comb. nov., 376; **cygnea** comb. nov., 376; **dispersa** comb. nov., 376; **erecta** comb. nov., 376, 378+; evaninervis, 374; **fendleri** comb. nov., 376; fontana, 369+*; f. var. pumila, 499; **glaziovii** comb. nov., 376; **globosa** comb. nov., 376; **le-testui** comb. nov., 376; **lindigii** comb. nov., 376; litorea, 176+*; longiseta, 462; **mexicana** comb. nov., 376; **nana** comb. nov., 376; **patula** comb. nov., 376; scabrifolia, 375, 383+; **sharpiana** sp. nov., 376+*; sp., 121; **tuerckheimii** comb. nov., 376; yezoana, 375
Philonotula curta, *376*; dregeana, 375; longiseta, 375; rigida, 375
Philophyllum, 270
Phylliscum demangeonii, 108
Physantholejeunea portoricensis, 331+
Physcia aipolia, 83+, 277; americana, 277; caesia, 108; millegrana, 277; phaea, 108; sp., 516; stellaris, 277; tenella, 83+
Physcomitrium coorgense, 184; pyriforme, 78+, 476
Physconia detersa, 277; distorta, 278; muscigena, 108, 502
phytogeography, of high-elevations in Mexico, 461+
Picea, 32; engelmannii, 105; glauca, 202
Pictolejeunea picta, 331+
Pilophorus pallidus, 69+

Pilopogon gracilis, 463; macrocarpus, 174; schilleri, 174+*; tiquipayae, 174
Pilotrichella flexilis, 463
Pilotrichidium, 270; callicostatum, 264*
Pinnatella alopecuroides, 184; ambigua, 184; intraimbata, 184
Pireella, 534; pohlii, 487
Placidiopsis pseudocinerea, 70
Placynthiella icmalea, 277; uliginosa, 277
Placynthium nigrum, 277
Plagiobryum demissum, 195, 476
Plagiochila abrupta, 330+; adiantoides, 330+*; arctica, 165; arcuata, 330+; asplenoides, 501; bicornis, 330+; bifaria, 330+; breuteliana, 330+; bursata, 330+; chinantlana, 330+; confundens, 330+; dichotoma, 330+; distinctifolia, 330+; divaricata, 330+; dominicensis, 330+; dubia, 330+; husnotii, 330+; hypnoides, 330+; jamaicensis, *347*; ludoviciana, 330+; micropterys, 330+; rutilans, 330+; simplex, 330+; sp., 121; superba, 330+; sylvicultrix, 330+; tenuis, 330+
Plagiomnium ellipticum, 480+, 499; medium, 480+, 499; m. var. **curvatulum** comb. nov., 533; rhynchophorum, 463
Plagiopus, 371+; oederi, 369+*; oederiana, 499
Plagiothecium, 159, 529; latebricola, 476; schraderi, 462
Platismatia glauca, 83+; tuckermanii, 277
Platydictya confervoides, new to Alabama, 487; jungermannioides, 499
Platygyriella densa, 462; pringlei, 462
Pleuridium ravenelii, new to Alablama, 487; subulatum, 487
Pleurochaete squarrosa, 464
Pleurophascaceae, 299
Pleurophascum grandiglobum, 301; var. decurrens, 301; **occidentale** sp. nov., 299+*
Pleuroziopsis, 159
Pleurozium, 529; schreberi, 203, 296, 499
Pogonatum, 323+; campylocarpum, 463; dentatum, 499; oligodus, 463; subflexuosum, 463
Pohlia andalusica, 476; annotina, 113; atropurpurea, new to Newfoundland, 476; cruda, 499; elongata, 464, 476; nutans, 112+, 296, 499; oerstediana, 462; wahlenbergii, 113; w. var. glacialis, 177
Polyblastia cucurbitula, 71
Polyblastiopsis, *71*; inductula, *71*; sublactea, *69*
Polycauliona phryganitis, *515*
Polychidium muscicola, 108+
Polysporina simplex, 277
Polytrichales, 324+
Polytrichastrum tenellum, 462
Polytrichum, 64+, 95, 397; alpinum, 499; commune, assessment of heavy metal toxicity on, 60+, 499; c. var. diminutum, 499; formosum, 195+; juniperinum, 464, 499; pallidisetum, new to Alabama, 483; piliferum, 499
Porella navicularis, 83+; swartziana, 330+
Porotrichodendron superbum, 463
Porotrichum cobanense, 463; longirostre, 463
Porpidia, 25+; albocaerulescens, 35+*, 277; **calcarea** sp. nov., 38+*; capillifolium, in North America, 25; **carlottiana** sp. nov., 39+*; cinereoatra, 39+*; contraponenda, 40+*; crustulata, 41+*, 108; **diversa** comb. nov., 42+*; **flavicunda** comb. nov., 43+*; flavocaerulescens, 108; flavocoerulescens, 43+*; glaucophaea, 47+*; **grisea** sp. nov., 48+*; **herteliana** sp. nov., 48+*; **lowiana** sp. nov., 49+*; macrocarpa, 50+*, 109, 277; musiva, 40+; platycarpoides, 69+; pseudomelinodes, 51+*; sp., 58; speirea, 51+*, 277; superba, 53+*; **tahawasiana** sp. nov., 53+*; **thomsonii** sp. nov., 54+*, 108+; trullisata, 33; tuberculosa, 55+*; zeoroides, 56+*
Pottia bryoides, 360; truncata, 112+; wilsonii, 357+*
Powellia, 268
predation, on bryophytes, 309
Preissa quadrata, 501
Prionodon densus, 464; dichotomus, 462; fusco-lutescens, 462
Prionolejeunea aemula, 331+; aequitexta, 345; exauriculata, 331+; helleri, 331+; innovata, 331+
proteins, a new method for separating and comparing, 164; separating and comparing in hepaticae, 164
protonemal growth, assessment of heavy metal toxicity on, 60; effect of hydrogen ion concentration on, 397+
Protoparmelia badia, 108
Protoparmeliopsis, 71; muralis, *69+*
Pseudephebe minuscula, 108; pubescens, 108, 502
Pseudobryum cinclidioides, 499
Pseudocrossidium **obtusulum** comb. nov., 533; revolutum, 356+, 361
Pseudocyphellaria anomala, 83+; anthraspis, 82+; crocata, 83+
Pseudoleskea arizonae, *395*; catenulata var. acuminata, 391; pallida, *390*; stenophylla, 473
Pseudoleskeella, 387+; arizonae, *395*; catenulata, 393+*; heteroptera, 387; nervosa, 388+*; n. var. sibirica, *390*; papillosa, 395+*, new to Newfoundland, 476, 499; **serpentinense** sp. nov., 391+*; **sibirica** comb. nov., 390+*; tectorum, 391+*
Pseudosymblepharis schimperiana, 463
Pseudotaxiphyllum, 159, 475
Psora decipiens, 502
Psoroma, 71; hypnorum, 502
Psorotichia schaereri, 277+
Pteryigynandrum catenulatum, 393; nervosum, 388
Pterobryaceae, 534
Pterobryopsis auriculata, 184; mexicana, 462
Pterogoniopsis, 529
Pterygoneurum kozlovii, 357+*; lamellatum, 360; ovatum, 356+, 360; subsessile, 357
Pterygophyllum, 262
Ptilidium ciliare, 501; pulcherrimum, 350+
Ptilium, 159; crista-castrensis, 499
Ptychocoleus polycarpus, *347*
Ptychomnion, 529
Punctelia rudecta, 455
PURSELL, R. A. Notes on neotropical Fissidens I, II, and III. I. The relationship of F. leptophyllus. II. The relationship of F. obtusissimus, stat. nov. III. The identity of F. hornschuchii, 523
Pycnolejeunea papulosa, 345; schwaneckei, 331+
Pylaisia, 210+; rhaphidostegioides, *210*
Pylaisiadelpha, 209+; brotherella, 210; drepanioides, 209; duellii, 210+; rhaphidostegioides, 209; sharpii, 210+; tenuirostris, 210+*; yokohamae, 210
Pylaisiella falcata, 464; intricata, 476; steerei, 419
Pyrenula nitida, 278; pseudobufonia, 455; santensis, 71
Pyrrhospora varians, 455

Quathlamba, 374+; debilicostata, 369+*, 374+
Quercus garryana, epiphyte succession on, 81+; rubra, distribution of corticolous lichens on, 453+

Racomitrium, 218; affine, 473; aquaticum, 195; canescens subsp. canescens, 476; crispulum, 464; elongatum, 473; ericoides, 476; heterostichum, 473; microcarpon, 473; steerei, 419; sudeticum, 473; venustum, 473
Racopilaceae, 268+
Racopilum, 268+; orthocarpum, 184; tomentosum, 464

Radula antillana, 330+; bolanderi, 82+; caldana, 344; complanata, 82+; dominicensis, *347*; evansii, 330+; fendleri, 344; flaccida, 330+; husnotii, 330+; inflexa, 344; kegelii, 330+; korthalsii, 344; laxiramea, 330+; longifolia, *344*, *347*; macrostachya, 330+; neotropica, 344; paganii, 330+; pallens, *347*; portoricensis, 344; prolifera, 501; recubans, 330+; saccatiloba, 330+; steerei, 419; subsimplex, 330+; surinamensis, *344*, *347*; tectiloba, 330+
RADWAN, S. S. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 178
Ramalina, 310+; americana, 277; calicaris, 278; complanata, 277; dilacerata, 455; farinacea, 82+, 277; fraxinea, 278; inflata, 82+; i. subsp. inflata, 71; intermedia, 277; pollinaria, 277; siliquosa, 310+
Ramonia valenzueliana, 70
Raphidorrhynchium lindigii, 462
Raphidostegium pylaisiadelphum, *210*
Rectolejeunea berteroana, 331+; brittoniae, 331+; emarginuliflora, 345; flagelliformis, 345; maxonii, 331+; phyllobola, 331+; pililoba, *347*
REDFEARN, JR., P. L., P.-C WU, S. HE & Y.-G. SU. Mosses new to mainland China, 183
REESE, W. D. Scopelophila cataractae in Texas, 413
REESE, W. D. Two new taxa of Syrrhopodon (Musci: Calymperaceae) from Australia, 302
REESE, W. D. *See* LIN, P.-J. & W. D. REESE, 170
REESE, W. D. & P.-J. LIN. Two new species of Syrrhopodon from Southeast Asia, 186
Regmatodon filiformis, 462
Regredicaulis monodactyla, *347*
Reinkella, 69; californica, *69*; parishii, 71; subcrustacea, *71*
Rhabdoweisia fugax, 464
Rhacocarpus purpurascens, 299
Rhacomitrium lanuginosum, 499
Rhaphidostegium, 210+
Rhizocarpon alpicola, 108+; badioatrum, 108; bolanderi, 106+; concentricum, 277; disporum, 108; eupetraeoides, 108+; eupetraeum, 106+; geminatum, 108+; geographicum, 108+; grande, 107+; macrosporum, 108; polycarpon, 107+; riparium, 108; superficiale, 106+
Rhizogoniaceae, 262+
Rhizogoniineae, 271
rhizoidal heads, in Leucobryum martianum, 493+
Rhizomnium, 161; andrewsianum, 499; appalachianum, 308, 479; glabrescens, 479; magnifolium, 308, 479; nudum, 479; pseudopunctatum, 308, 479+; punctatum subsp. chlorophyllosum, 308; p. var. punctatum, 479
Rhizoplaca chrysoleuca, 108; melanopthalma, 108
Rhodobryum roseum, 464
Rhododendron campanulatum, 122
Rhynchostegiopsis, 270; carolae, 266; flexuosa, 266, 463
Rhytidadelphus squarrosus, 308
Rhytidium, 159; rugosum, 499
Riccardia digitiloba, 331+; fucoidea, 331+*; fucoides, *347*; latifrons, 350+; palmata, 350+; pinguis, *347*; schwaneckei, 331+; sprucei, 331+
Riccia, 120; brittonii, 346; elliottii, 331+; membranacea, 331+; violacea, 331+
Rigodium, 159; gracile, 462
Rimularia furvella, 69+; impavida, 69+; insularis, 71
Rinodina calcigena, 71+; endophragmia, 71; exigua, 277; occidentalis, *71*+; sophodes, 278; turfacea, 108
ROBINSON, A. L., D. H. VITT & K. P. TIMONEY. Patterns of community structure and morphology of bryophytes and lichens relative to edaphic gradients in the subarctic forest-tundra of northwestern Canada, 495
Rozea subjulacea, 462
RUSHING, A. E. Bruchia brevifolia new to Alabama, 130; Bruchia brevifolia Sull., 260
RYAN, B. D. A monograph of Lecanora subg. Placodium sect. Endochloris (lichenized Ascomycotina), 513

Saccharum spontaneum, 121
Saccogyna, 123
SALAZAR A., N. A preliminary report of rhizoidal heads in Leucobryum martianum, 493
Sanionia uncinata var. **symmetrica** comb. nov., 533
Sasaokaea, 162
Sauloma, 270; macrospora, 266; tenella, 176
Scapania degenii, 501; gymnostomophila, 501; irrigua, 501; mucronata, 501; nemorosa, 164; paludicola, 501; paludosa, 501; portoricensis, 330+; simmonsii, 501; sp., 121
Schaereria cinereorufa, 109; tenebrosa, 109+
Schiffneriolejeunea polycarpa, 330+
Schimperobryum, 270; splendissimum, 266
Schismatomma rappii, 71
Schistidium maritimum, 475; rivulare var. **latifolium** comb. nov., 533; steerei, 419
Schistoplaca, 69
Schizomitrium, 270; diatomophilum, 266; pallidum, 266, new to Alabama, 487
Schlotheimia aciculare, 282; baileyi, 283+, 290; brownii, 290+*; campbelliana, 283+*; cirrosa, 282; densifolia, 283; funiformis, 293+*; knightii, 286+*; quadrifida, 282+; rhystophylla, 283; rugifolia, 282+; squarrosa, 282+; sulcata, 282; torquata, 282+; viticulosa, 282
Schoenobryum concavifolium, 184
SCHOFIELD, W. B. *See* ANDO, H., T. SEKI & W. B. SCHOFIELD, 209
Sclerophora nivea, 277
Scoliciosporum chlorococcum, 277+; umbrinum, 277
Scopelophila cataractae, in Pennsylvania, 112+; in Texas, 413+; ligulata, 112
Scorpidium scorpioides, 161, 195, 499; turgescens, 195, 480, 499
SEKI, T. *See* ANDO, H., T. SEKI & W. B. SCHOFIELD, 209
Seligeria brevifolia, 476
Sematophyllaceae, 529
Sematophyllum, 210; homomallum, 299; sericifolium, 463
SHARP, A. J. [Review of:] A. Noguchi, Illustrated Moss Flora of Japan. Part 2, 555
SHAW, J. & S. C. BEER. Scopelophila cataractae (Mitt.) Broth. in Pennsylvania, 112
SHAW, J., S. C. BEER & J. LUTZ. Potential for the evolution of heavy metal tolerance in Bryum argenteum, a moss. I. Variation within and among populations, 73
SHAW, J., B. D. MISHLER & L. E. ANDERSON. Peristome development in mosses in relation to systematics and evolution. IV. Haplolepideae: Ditrichaceae and Dicranaceae, 314
SNIDER, J. A. [Review of:] H. Crum in collaboration with S. Planesek, A Focus on Peatlands and Peat Mosses, 256
SÖDERSTRÖM, L. Regional distribution patterns of bryophyte species on spruce logs in northern Sweden, 349
Solenostoma crenulata, 74
Solorina, 203+; bispora, 502; crocea, 203, 502
South Africa, weathering of rock by lichens in, 407+
South America, Lejeunea capensis, disjunct with Africa, 305
Sphaerophorus fragilis, 502; globosus, 502
Sphaerotheciella pinnata, 462
Sphagnoecetis portoricensis, *344*
Sphagnum, 1+, new tropical American species of, 98+, 181, 194+, 324+, 397+; sect. Acutifolia, 20+; sect.

Cuspidata, 20+; sect. Rigida, 20; sect. Sphagnum, 20+; sect. Squarrosa, 20+; sect. Subsecunda, 20; alegrense, 405; andersonianum, 3+*; angustifolium, 4, 117, 499; aongstroemii, 499; apollinairei, 100; **aureum** sp. nov., 405+*; balticum, 499; bartlettianum, 488; capillifolium, biosystematic study of, 1+*, 100, 499; c. var. capillifolium, 19+*; c. var. tenellum, 1+*; c. var. tenerum, 1; centrale, 4; **cleefii** sp. nov., 100+*; compactum, 196; contortum, 196, 499; cuspidatum, 103; cyclophyllum, 102; fallax, 117+; fimbriatum, 499; fitzgeraldii, 19; flexuosum, 117; fuscum, 4, 499; girgensohnii, 4, 499; **imperforatum** sp. nov., 98+*; lescurii, 20; limbatum, 463; magellanicum, 3+, 181, 196, 405, 499; mendocinum, 119, 195; meridense, 405; nemoreum, 6+*; obtusum, 500; **pacificum** sp. nov., 116+; perichaetiale, 98, 405; **poasense** sp. nov., 102+*, pulchricoma, 117; recurvum, 117; recurvum complex, 116+; richardsianum, 104; riparium, 196; rubellum, 2+*; russowii, 4, 500; sparsum, 100+, 405; squarrosum, 500; steerei, 419; **subdivisum** sp. nov., 98+*; subtile, 2+*; teres, 500; **trirameum** sp. nov., 104+*; warnstorfii, 500

Sphinctrina sp., 519; turbinata, 455

Spilonema revertens, 106, 107+; tuberculosa, *55*

Splachnaceae, 397+

Splachnobryum obtusum, new to Alabama, 488

Splachnum, neutrophilic hypothesis of substrate restriction in, 397+; ampullaceum, 397; luteum, 398+; sphaericum, 398+

Sporastatia polyspora, 109; testudinea, 106+

spore germination, assessment of heavy metals toxicity on, 60; effect of hydrogen ion concentration on, 387+

Squamidium, 268

SRIVASTAVA, S. C. & A. K. ASTHANA. Two species of Folioceros from India, including F. kashyapii, sp. nov., 219

Staurothele clopima, 107+; discedens, 71; succedens, 71

Steerea, 418; mastigophoroides, 418

Steerecleus, 418; serrulatus, 418

Steereella, 418; linearis, 418

Steereobryon, 418; subulirostrum, 418

Steereochila, 418; ecuadorica, 418

Steereocolea, 418; bisbifida, 418

Steereomitrium, 418; minutum, 418

Stegonia latifolia var. pilifera, 500; **pilfera** comb. nov., 533

Stenodesmus, 268+

Stenodictyon, 265; nitidum, 266

Stephanocyclos, 31; hensennianus, 33

Stereocaulon alpinum, 502; dactylophyllum, 278; glareosum, 502; paschale, 203+, 502; saxatile, 277; tomentosum, 203, 277, 502

Stereodon, 209+; subg. Heterophyllium, 209+; subg. Pseudo-Rhaphidostegium, 209; lorentzianus, 209; nemorosus, 209+; yokohamae, *210*

Sticta fuliginosa, 83+; limbata, 83+

Stictolejeunea squamata, 330+

Stigonema, 25, 110

STONE, D. F. Epiphyte succession on Quercus garryana branches in the Willamette Valley of western Oregon, 81

STONEBURNER, A. *See* WYATT, R. & A. STONEBURNER, 299, 308

STONEBURNER, A. & R. WYATT. Notes on Rhizomnium pseudopunctatum and associated Mniaceae from central New York State, 479

STOTLER, R. E. & B. CRANDALL-STOTLER. Recent Literature on Hepatics and Anthocerotes—117, 131;—118, 228;—119, 420;—120, 535

Strangospora moriformis, 277

Strepsilejeunea involuta, *347*

Streptopogon erythrodontus, 464

STUBBS, C. S. Patterns of distribution and abundance of corticolous lichens and their invertebrate associates on Quercus rubra in Maine, 453

Stylolejeunea pililoba, *347*

SU, Y.-G. *See* REDFEARN, JR., P. L., P.-C. WU, S. HE & Y.-G. SU, 183

Sulfur dioxide, lichen sensitivity to, 275+

Sweden, distribution patterns of bryophytes in, 349

Sykoraea, nom. nud., 123

Sykorea, 123; viticulosa, 123

Symbiezidium barbiflorum, 330+; granulatum, *347*; laceratum, *347*; transversale, 330+; vincentinum, *347*

Symblepharis vaginata, 464

Symphydon echinatus, 267

Symphyogyna aspera, 331+; brongniartii, 331+; trivittata, 331+

Syringothecium, *529*+; brasiliense, *531*; nemodontium, *529*; sprucei, *529*

Syrrhopodon, 170+, new species in China, 186+; subg. Syrrhopodon, 303; albidus, 304; armatispinosus, 170+; armatus, 170+; borneensis, 170+; **chenii** sp. nov., 170+, 186+*; flammeonervis, 170+; gardneri, 170+; gaudichaudii, 187; involutus, 170+; japonicus, 170+; konoi, 170+; larminatii, 170+; lonchophyllus, 170+; **orientalis** sp. nov., 170+, 186+*; parasiticus var. parasiticus, 170+; prolifer, 170+; p. var. **mossmanensis** var. nov., 303+*; p. var. prolifer, 304; prolifer complex, 304; semperi, 170+; spiculosus, 170+; **stoneae** sp. nov., 302+*; tjibodensis, 170+; tosaensis, 170+; trachyphyllus, 170+

Syzygiella laevigata, *347*; perfoliata, 330+; rubricaulis, *347*

Taxilejeunea antillana, 346; eggersiana, 346; obtusangula, 331+; pterigonia, 331+, 345; sulphurea, 331+

Taxiphyllum, 159; taxirameum, 483

Tayloria acuminata, 500; froelichiana, 500; mexicana, 462

Telaranea nematodes, 330+; sejuncta, *347*

Tephromela, 71; atra, 519

Tetraphis pelucida, 323, 350+

Tetraplodon, 196; angustatus, 195; mnioides, 397, 500; pallidus, 500; paradoxus, 500

Tetrastichium, 267; fontanum, 266

Tetrodontium repandum, 476

Texas, Scopelophila cataractae new to, 413+

Thamniopsis, 270; pendula, 266

Thamnobryum, 159

Thamnolia subuliformis, 467+, 502; vermicularis, 467+, 502

Thelenella, 71; harrisii, 71; hassei, 70+; inductula, 70+; modesta, 70+; sordidula, 70+; weberi, 71

Thelidium parvulum, 277

Thelomma sp., 521

THOMSON, J. W. Additions and a revised key to Catapyrenium in North America, 190

Thuidiopsis furfurosa, 462

Thuidium, 96; abietinum, 500; delicatulum, 121; d. var. peruvianum, 462; d. var. radicans, 464; mexicanum, 464; minutulum, 464, 488; recognitum, 500; tamariscellum, 121+

Tierra del Fuego, additions to the moss flora of, 174

TIMME, S. L. *See* BOWERS, F. D., R. R. HAYNES & S. L. TIMME, 482

Timmia sect. Timmiaurea, 371+; austriaca, 369+*, 500; megapolitana, 369+*, 500; norvegica, 369+*, 500

Timmiaceae, 368+

TIMONEY, K. P. *See* A. L. ROBINSON, D. H. VITT & K. P. TIMONEY, 495

Tomasellia sparsella, 277
Tomenthypnum, 500
Tomentypnum, 158; nitens, 500
Tortella fragilis, 500; tortuosa, 500
Tortula, 112; amphidiacea, 462; bistratosa, 360; brevipes, 360; caninervis, 360; fragilis, 464; inermis, 411+*; norvegica, 500; pagorum, 483; rhizophylla, new to Alabama, 483; ruralis, 356+, 500; scotteri, 360
toxicity, assessment of heavy metal combinations, 60
Trachylejeunea aquarius, 331+; inflexa, *347*
Trachypodopsis, 120; crispatula, 121+
Trapelia coarctata, 277; involuta, 277; placodioides, 277
Trapeliopsis flexuosa, 277; granulosa, 277; viridescens, 277
Trebouxia, 25
Trematodon longicollis, 314+, 488
Trichocolea brevifissa, 329+; elliottii, 344; filicaulis, 329+; flaccida, 329+; tomentosa, 329+
Trichodon cylindricus, 500
Trichostomopsis australasiae, 360
Trichostomum arcticum, 473; brachydontium, 464; crispulum, new to Alabama, 483; spirale, *533*
Triquetrella curvifolia, *383*+; tasmanica, 383+
Tritomaria exectiformis, 501; polita, 501; quinquedentata, 501; scitula, 501
tropical America, new species of Sphagnum from, 98
Trypethelium virens, 278
TUCKER, S. C. [Review of:] I. M. Brodo, Lichens of the Ottawa Region. Second Edition, 257
Tuckermannopsis chlorophylla, 83+; oakesiana, 455; pinastri, 109
Tylimanthus marginatus, 330+

Ulota, 217
Umbilicaria cylindrica, 109; deusta, 109; hyperborea, 109; lambii, 106+; phaea, 109; polyphylla, 109; proboscidea, 109; rigida, 109; torrefacta, 107+; vellea, 106+
United States, changes to the lichen checklist of, 68
Urceolaria actinostoma, *69*; albissima, *69*+
Usnea, 82+; barbata, 278; caucasica, *71*; cornuta, 71+; dasypoga, *71*; esthonica, *71*; filipendula, 71, 82+; flagellata, *71*; fragilescens var. mollis, 71+; fulvoreagens, 71; hirta, 278; inflata, *71*+; lapponica, 71; laracina, *71*; longissima, 82+; michauxii, 71; mollis, *71*+; monstruosa, 71; perplexans, 71; rubiginea, *71*; sorediifera, *71*; strigosa, 71; s. subsp. major, 71; s. subsp. rubiginea, 71; s. subsp. strigosa, 71; subfloridana, 82+, 455; substerilis, 71

Verbascum nigrum var. bracteatum, 123; sykoraeum, 123
Verrucaria fuscella, 278; muralis, 109, 278; mutabilis, 71; nigrescens, 278; viridula, 278
Vezdaea stipitata, 82+
Viola calaminaria, 112
VITAL, D. M. *See* GIANCOTTI, C. & D. M. VITAL, 305
VITT, D. H. The genus Schlotheimia (Orthotrichaceae: Bryopsida) in Australia and New Zealand, 282
VITT, D. H. *See* A. L. ROBINSON, D. H. VITT & K. P. TIMONEY, 495

WALKER, M. D. Microhabitat preferences of Dactylina arctica and D. beringica on pingos in northern Alaska, 467
Warnstorfia, 161; **crassicostata** comb. nov., 533; fluitans var. **falcata** comb. nov., 533
Washington, Didymodon vinealis var. luridus new to, 361; Pseudocrossidium revolutum new to, 361; Trichostomopsis australasiae new to, 360
weathering, of rock by Acarospora sinopica, 407+
Weissia, 411; bartramioides, *376*; brachycarpa, 361; controversa, 74+, 225, 464, 500; jamaicensis, 463, 488
WERGER, M. J. A. & H. J. DURING. A subterranean moss greenhouse in the Californian desert, 411
Weymouthia, 268
WHITTEMORE, A. [Review of:] W. R. Buck (ed.), Bryostephanae Steereana: A Collection of Bryological Papers Presented to William Campbell Steere on the Occasion of his 80th Birthday, 153
WHITTEMORE, A. [Review of:] R. C. Kruijt, A Monograph of the Genera Dicranolejeunea and Acanthocoleus, 448
WHITTEMORE, A. & B. ALLEN. The systematic position of Adelothecium Mitt. and the familial classification of the Hookeriales (Musci), 261
Wijkia costaricensis, 462
WILSON, P. & D. H. NORRIS. Pseudoleskeella in North America and Europe, 387
WU, P.-C. *See* REDFEARN, JR., P. L., P.-C. WU, S. HE & Y.-G. SU, 183
WYATT, R. *See* CAMERON, R. G. & R. WYATT, 397
WYATT, R. *See* STONEBURNER, A. & R. WYATT, 479
WYATT, R. & A. STONEBURNER. Pleurophascum occidentale: A new moss from Western Australia, 299
WYATT, R. & A. Stoneburner. Bryophytophagy of Rhizomnium punctatum by larvae of the crane fly Tipula oropezoides, 308

Xanthoparmelia australasica, 71; centrifuga, 502; congensis, *71*+; cumberlandia, 326+; mougeotina, 70+; neochlorochroa, 71; norchlorochroa, 71; norhypopsila, 71; separata, 502; somloensis, 109; stenophylloides, 71
Xanthoria candelaria, 109, 516; elegans, 107+; fallax, 278; polycarpa, 278

ZAVADA, M. S. *See* KERR, S. & M. S. ZAVADA, 407
Zoopsidella monodactyla, 330+
Zoopsis antillana, 330+; monodactyla, *347*
Zygodon, 174+; hookeri, 176; magellanicus, 176; menziesii, 174+*; obtusifolius, 176; pichinchensis, 176; reinwardtii, 464; r. var. hyadesii, 176; viridissimus, 464